

Rabbit Anti-D₃ Receptor Sera

AS-3532S

Lot # 7851

The antiserum was raised in a rabbit which was immunized with D₃(2-10-Cys¹¹) covalently attached onto a carrier protein, and it has been characterized by immunocytochemical, western immunoblot and ELISA techniques. This antiserum has been found to be highly specific for this peptide sequence and is suitable for immunocytochemical detection of the D₃ dopamine receptor. Rehydrate the lyophilized antibody with 0.1 ml of PBS which contains 10 mg/ml BSA or with additional buffer (such as 10 mg/ml BSA in PBS) for more dilute antibody.

Antiserum Specificity

Polypeptide	% Cross Reactivity
D ₃ Dopamine Receptor(2-10)	100
D ₃ Dopamine Receptor	80
D ₁ Dopamine Receptor(9-21)	0
D ₁ Dopamine Receptor	0
D ₂ Dopamine Receptor(272-282)	0
D ₂ Dopamine Receptor	0
D ₄ Dopamine Receptor(176-185)	0
D ₄ Dopamine Receptor	0
D ₅ Dopamine Receptor(23-35)	0
D ₅ Dopamine Receptor	0

Immunocytochemical Staining

This antiserum has been found to stain specific cells in various regions of PLP fixed rat brain sections at a dilution of 1:5000. This includes dendritic labeling in the molecular layers (cortex layer 1 and molecular layer of the hippocampus) as well as the plasma membranes of neurons in layer 5 of cortex and the glomeruli of the olfactory bulb. For additional information on PLP fixative see the FAQ page on our web site at www.RDAbs.com.

Western Immunoblot

Western immunoblots using whole rat brain homogenate resulted in a single band being detected at ~50 KDa at 1:500 dilution.

Western Blotting Protocol

1. After SDS-PAGE (on either 4-15% gradient gels or single percentage gels, such as 12% gels) and electrophoretic transfer to PVDF membrane, block the membrane overnight with 2% normal goat serum in TBS/Tween-20 buffer.
2. Wash x 2 with TBS/Tween-20.
3. Apply the rabbit polyclonal antibody after dilution to at least 1:500 (Note: higher dilutions may be needed). Use 2% normal goat serum in TBS/Tween-20 as buffer for the primary antibody. Let the primary antibody bind for 2-4 hours.
4. Wash x 3 with TBS/Tween-20.
5. Apply affinity purified HRP-goat anti-rabbit IgG antiserum diluted 1:2500 (dilution may vary depending upon supplier) in 2% normal goat serum in TBS/Tween-20. Incubate 1-2 hours. Note: greater sensitivity may be achieved using ABC techniques.
6. Wash x 4 for 5 minute/wash with TBS/Tween-20.
7. Develop color using the enhanced DAB reaction.